

Oscillations in the position of the maximum of a lattice branching random walk and application to the binary search tree

Pierre-Antoine Corre

November 18, 2014

Abstract

For a particular case of branching random walk with lattice support, namely the Yule process, we prove that the distribution of the centred maximum oscillates around a distribution corresponding to a critical travelling wave in the following sense: there exist continuous functions $t \mapsto a_t$ and $x \mapsto \phi_{\max}(x)$ such that:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(X_{\max}(t) \leq a_t + x) - \phi_{\max}(x - \{a_t + x\})| = 0,$$

where $\{x\} = x - \lfloor x \rfloor$. This process is classically related to the binary search tree, thus yielding similar results for the height and for the saturation level of the binary search tree too.

1 Introduction

1.1 Main results

We consider two models of growing binary trees: the binary search tree $(\mathcal{T}_n)_{n \in \mathbb{N}^*}$ and the Yule tree process $(X_t, t \geq 0)$. The binary search tree is a Markov chain on the space of binary trees that can be constructed as follows: $\mathcal{T}_1$ is a tree made of a single leaf. To obtain $\mathcal{T}_{n+1}$ we uniformly choose a leaf of $\mathcal{T}_n$ and attach to it two leaves. For more details and others constructions see e.g. [10, 28].

The Yule tree process is a growing tree which can be seen as a continuous time version of $(\mathcal{T}_n)_{n \in \mathbb{N}^*}$. Specifically, at time 0 the tree is reduced to a leaf. Each leaf lives for a random time with exponential β distribution independent of the others leaves. When it dies, it is replaced by two children. Moreover, we give a spatial structure to the Yule tree process. Let $\mathcal{N}_t$ be the set of leaves alive at time t and N_t its cardinal, for $u \in \mathcal{N}_t$, the position $X_u(t)$ of u at time t is its generation. For $s \leq t$, if $v \in \mathcal{N}_s$ is an ancestor of $u \in \mathcal{N}_t$ then $X_u(s) = X_v(s)$ by definition. The process with spatial structure, which we will call the Yule-time process as in [8] is then a branching random walk with continuous time and lattice-integer support.

Note that N_t jumps by increments of 1 and that if we stop a Yule tree process at time

$$\tau_n = \inf\{t > 0, N_t = n\}, \tag{1}$$

we have:

$$(\mathcal{T}_n)_{n \in \mathbb{N}} \stackrel{\mathcal{L}}{=} (X_{\tau_n})_{n \in \mathbb{N}}. \quad (2)$$

Our goal is to study the asymptotic behaviour of the extremal particles of the Yule-time process and to deduce from it the asymptotic behaviour of the height (i.e. the highest generation of a leaf) and of the saturation level (i.e. the maximal level h such that there are no leaves for all levels up to h) of the binary search trees.

Since Bramson [6], we know exactly what happens for another spatial branching process: the branching Brownian motion (BBM). If $X_{\max}(t)$ is the maximal position of a particle in the BBM and m_t is its median then $p(t, x) = \mathbb{P}(X_{\max}(t) \leq x)$ solves the Kolmogorov-Petrovskii-Piskunov equation (KPP) [21] and

$$\lim_{t \rightarrow +\infty} p(t, x + m_t) = \phi(x), \quad (3)$$

where $\phi \in \mathcal{C}^1(\mathbb{R})$ is a critical travelling-wave solution of KPP. By travelling-wave solutions we mean a solution $h \in \mathcal{C}^1(\mathbb{R} \times \mathbb{R}^+)$ of the form $h(x, t) = F(x - ct)$ with $(F, c) \in \mathcal{C}^1(\mathbb{R}) \times \mathbb{R}$ (c is called the speed of the travelling-wave) and by critical we mean a travelling-wave solution at speed $c^* = \lim_{t \rightarrow +\infty} X_{\max}(t)/t$. The travelling-wave solution at the critical speed is unique in the class of monotone bounded functions solutions up to a translation in the argument.

Let us now consider the Yule process and define $X_{\max}(t)$ and $X_{\min}(t)$ to be the maximal and minimal position of it. For $(x, t) \in \mathbb{R} \times \mathbb{R}^+$:

$$\begin{cases} h_{\max}(x, t) = \mathbb{P}(X_{\max}(t) \leq x) = \mathbb{P}(X_{\max}(t) \leq \lfloor x \rfloor) \\ h_{\min}(x, t) = \mathbb{P}(X_{\min}(t) \geq x) = \mathbb{P}(X_{\min}(t) \geq \lceil x \rceil) \end{cases}. \quad (4)$$

Both solve the equation:

$$\partial_t h(x, t) = \beta(h^2(x - 1, t) - h(x, t)), (x, t) \in \mathbb{R} \times \mathbb{R}^+. \quad (5)$$

Note that equation (5) is the analogue of the KPP equation. One can easily check that ϕ is a travelling-wave solution of (5) with speed c if and only if it solves the differential equation:

$$c\phi'(x) = \beta(\phi(x) - \phi^2(x - 1)), \quad x \in \mathbb{R}. \quad (6)$$

Let us recall an existence and uniqueness result for (6). First, introduce the speeds of the extremal particles of the Yule-time process:

$$\lim_{t \rightarrow +\infty} \frac{X_{\max}(t)}{t} = c^+ \quad \text{and} \quad \lim_{t \rightarrow +\infty} \frac{X_{\min}(t)}{t} = c^-. \quad (7)$$

For $\theta \in \mathbb{R}^*$, let

$$c_\theta = \frac{(2e^\theta - 1)\beta}{\theta}. \quad (8)$$

Then $\theta^+ = 0.768\dots$ is the largest solution of $c_\theta = 2e^\theta\beta$ and $\theta^- = -1.678\dots$ is the smallest. We have $c^- = c_{\theta^-} = \beta \times 0.373\dots$ and $c^+ = c_{\theta^+} = \beta \times 4.311\dots$

Theorem (Chauvin, Rouault [8]). *Equation (6) has monotone and bounded travelling-wave solutions at speed c , in $\mathcal{C}^1(\mathbb{R})$ if and only if $c \leq c^-$ or $c \geq c^+$. Moreover, uniqueness holds for each such $c \neq 0$ (up to an additive constant in the argument). For $c = 0$, the whole set of solutions is:*

$$\{x \mapsto P(x)^{2^x} \mid P \text{ is 1-periodic and nonnegative}\}. \quad (9)$$

It is easy to see that we can find more than one monotone bounded solution in the set (9).

Let us point out that this result is stated in a slightly different form in [8]. Indeed, Chauvin and Rouault proved uniqueness for this equation up to a decreasing change of variables and in the class of Laplace transforms. The key to their proof was an application of Liu [24, 25]. The more recent result from Alsmeyer, Biggins and Meiners [4] gives us the uniqueness among decreasing functions in $[0, 1]$. We observe that the set of monotone bounded solutions of (6) is in fact the set of monotone solutions in $[0, 1]$. As far as non-existence of solutions at other speeds is concerned note that we can exactly transpose the proof from Harris [19] for the branching Brownian motion.

Let us now consider the asymptotic distributions of the extremal particles. Aïdekon [2] proved a result similar to [6] for a large class of branching random walk in the non-lattice case. Bramson, Ding and Zeitouni [7] have recently given an alternative proof for the non-lattice case in a slightly less general case. In particular, they assume that the displacements of each offspring of a particle are independent. Furthermore, the authors suggest that their method works in the lattice case. The Yule-time process does not satisfy the assumptions of [7] but the result will be the same. The Yule-time process evolving on a lattice support, the functions $h_{\max}$ and $h_{\min}$ defined in (4) are continuous in t but piecewise constant in x , which implies that whatever the centring is, the distribution cannot converge. However, we will see in the following theorem that the asymptotic distributions oscillate around the critical travelling-wave.

Theorem 1.1. *Let $a_t = c^+t - \frac{3\log(t)}{2\theta^+}$, $b_t = c^-t - \frac{3\log(t)}{2\theta^-}$ (recall that $\theta^- < 0$) and $\{x\} = x - \lfloor x \rfloor$.*

There exists a travelling-wave solution $\phi_{\max}$ at speed c^+ and a solution $\phi_{\min}$ at speed c^- of (5) such that:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(X_{\max}(t) \leq a_t + x) - \phi_{\max}(x - \{a_t + x\})| = 0 \quad (10)$$

and:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(X_{\min}(t) \geq b_t + x) - \phi_{\min}(x - \{b_t + x\})| = 0. \quad (11)$$

To study the maximum of the Yule-time process we change our point of view and we introduce the Yule-generation process defined as follow. Let $u \in \mathcal{M}_n$, where $\mathcal{M}_n$ is the set of the particles of the n th generation, we define

$$T_u(n) = \inf\{t > 0, X_u(t) = n\} \text{ and } (T_n) = \{T_u(n), u \in \mathcal{M}_n\}. \quad (12)$$

It is clear that (T_n) is simply the branching random walk X stopped on successive stopping lines $(X = n)$ and that (T_n) is obtained from X switching time and space. Furthermore, (T_n) itself is a branching random walk with discrete time and continuous spatial position with the following branching mechanism. At time $n = 0$, a particle is at 0. At time $n = 1$ the particle dies and gives birth to the point process: $\xi = 2\delta_T$ where T is distributed as an exponential variable with parameter β . At each time n , the particles of the previous generation die and give birth to particles whose displacements from the parents are given by

i.i.d. copies of ξ .

This connection was established by Chauvin and Rouault (2005) [8], which allowed them to prove the existence and uniqueness of travelling-waves.

An advantage of such a change of point of view is that the non-lattice case is better understood and that the results in this framework can be transposed thanks to the relation:

$$\mathbb{P}(T_{\min}(n) \leq t) = \mathbb{P}(X_{\max}(t) \geq n), \quad (13)$$

where $T_{\min}(n) = \inf\{T_u(n), u \in \mathcal{M}_n\}$. Indeed, the event $\{T_{\min}(n) \leq t\}$ means that the first apparition time of a particle of generation n is before t , which is equivalent to the fact that the maximal generation of a particle at time t is greater than n . We give below an illustration of this correspondence. The generation of the particle u in red at time $t = 4$ is 2 and the reaching time of the second generation for u is $T_u(2)$. By switching time and generation we can see that the Yule generation process is a branching random walk with discrete time and non-lattice support.




Figure 1: Correspondence between Yule-time process and Yule generation process

The Yule generation process being a non-lattice branching random walk Aidekon's result [2] applies. Therefore, if the number of children of a particle is a random variable N and if the life-time has a distribution G which satisfies the assumptions of Aidekon's result, Theorem 1.1 holds.

We will now extend Theorem 1.1. For each piecewise continuous function f , we define H_f and P_f by:

$$H_f(t) := \mathbb{E}[f(X_{\max}(t) - a_t)], \quad t > 0 \quad (14)$$

and:

$$P_f(s) := \sum_{k \in \mathbb{Z}} f(k - s) (\phi_{\max}(k - s) - \phi_{\max}(k - 1 - s)), \quad s \in \mathbb{R} \quad (15)$$

when these functions are well-defined. Note that P_f is 1-periodic. Taking, for $x \in \mathbb{R}$, $f(y) = \mathbf{1}_{\{y \leq x\}}$, Theorem 1.1 can be seen (if we omit the uniformity of the convergence) as a particular case of the following corollary.

Corollary 1.2. *There exists $\delta > 0$ such that for all piecewise continuous function f satisfying $f(x) = o_{\pm\infty}(e^{\delta|x|})$ we have:*

$$\lim_{t \rightarrow +\infty} |H_f(t) - P_f(a_t)| = 0. \quad (16)$$

In particular, this yields for $f = \text{Id}$ that there exists a 1-periodic function P such that:

$$\lim_{t \rightarrow +\infty} |\mathbb{E}(X_{\max}(t)) - a_t - P_f(a_t)| = 0. \quad (17)$$

The analogue corollary for the minimum of the process also holds. The case where $f = \text{Id}$ is of particular interest. Indeed, by differentiating u , $u(t) := \mathbb{E}(X_{\max}(t))$, we obtain that $u'(t) = \mathbb{E}(F_t)$, where:

$$F_t = \text{Card}\{u \in \mathcal{N}_t, X_u(t) = X_{\max}(t)\}. \quad (18)$$

Drmotá [16] proved that if we call C_n the number of particles on the highest generation in a binary search tree with n leaves, then there exists a 1-periodic function P such that $\mathbb{E}(C_n) - P(a_{\log n})$ converges to 0. Corollary 1.2 is not precise enough to prove a similar result. The following corollary gives us more accuracy on the derivatives of P_f and H_f .

Corollary 1.3. *Let δ be defined as in the previous corollary. Let $f \in \mathcal{C}^k(\mathbb{R})$ satisfying $f^{(i)}(x) = o_{\pm\infty}(e^{\delta|x|})$, for $0 \leq i \leq k$.*

Let $g(t) = P_f(a_t)$. We have that $H_f, g \in \mathcal{C}^k(\mathbb{R}^+)$ and:

$$\lim_{t \rightarrow +\infty} |H_f^{(k)}(t) - g^{(k)}(t)| = \lim_{t \rightarrow +\infty} |H_f^{(k)}(t) - (-c^+)^k P_f^{(k)}(a_t)| = 0. \quad (19)$$

In particular, this yields for $f = \text{Id}$ that there exists a 1-periodic smooth function P such that:

$$\lim_{t \rightarrow +\infty} |\mathbb{E}(F_t) - P(a_t)| = 0 \quad (20)$$

Whether P in the former corollary is or is not constant is still an open question. We discuss this issue in more detail in the next section.

Knowing the asymptotic distribution of the extremal particles of the Yule tree process, we can obtain the corresponding analogue result for the asymptotic distributions of the height and the saturation level of the binary search tree, by

using the connection between the two models. We call $\partial M_\infty(\theta^+)$ and $\partial M_\infty(\theta^-)$ the limit of the derivative martingales of the binary search tree defined in (65), and for $C > 0$:

$$\psi_C^+(x) = \mathbb{E} \left(e^{-C e^{-\theta^+ x} \partial M_\infty(\theta^+)} \right) \text{ and } \psi_C^-(x) = \mathbb{E} \left(e^{-C e^{-\theta^- x} \partial M_\infty(\theta^-)} \right). \quad (21)$$

Theorem 1.4. *Let H_n be the height (i.e. the highest generation of a leaf) of a random binary search tree with n nodes and h_n its saturation level (i.e. the maximal level h such that there are no leaves for all levels up to h). There exist $C^+, C^- > 0$ such that:*

$$\lim_{n \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) - \psi_{C^+}^+(x - \{a_{\log(n)} + x\})| = 0, \quad (22)$$

and:

$$\lim_{n \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(h_n \geq \lceil b_{\log(n)} + x \rceil) - \psi_{C^-}^-(x - \{b_{\log(n)} + x\})| = 0. \quad (23)$$

This result can be generalised like Theorem 1.1. If the offspring distribution of a leaf, denoted by N , satisfies $\mathbb{E}(N \log^2 N) < +\infty$, then Theorem 1.4 holds.

1.2 Discussion

Several approaches have been used in the study of (5). Drmota [14], proved that there is a unique solution to the delay differential equation:

$$\Phi'(x) = -\frac{1}{\alpha^2} \Phi\left(\frac{x}{\alpha}\right)^2, \quad (24)$$

where $\alpha = e^{\frac{\beta}{c^+}}$, $\Phi \in \mathcal{C}^1(\mathbb{R}^+, \mathbb{R})$ and with initial condition $\Phi(0) = 1$. We can check that if

$$f(x) = e^{\frac{\beta x}{c^+}} \Phi\left(e^{\frac{\beta x}{c^+}}\right), \quad (25)$$

where Φ is a solution of (24), then f is a solution of (6) with $c = c^+$. Moreover he found the following asymptotic for Φ :

$$1 - x\Phi(x) \underset{x \rightarrow +\infty}{\sim} C \frac{\log x}{x^{c^+-1}} \quad (26)$$

with $C > 0$. Recall that Chauvin and Rouault prove that solutions of (6) are unique (up to a shift). Their method also gives the uniqueness of solutions of (6) for a slightly different class of functions, it allows us to determine the left tail of the travelling-waves ϕ_c with speed $c \geq c^+$ of (5):

$$\phi_c(x) \underset{x \rightarrow -\infty}{\sim} e^{\frac{\beta x}{c}} \quad (27)$$

Let us now mention some related works to Theorem 1.1. Lifshits in [23] considered the following branching random walk. At time $n = 0$ a particle is at 0. At each time $n \in \mathbb{N}$ every particle produces two particles which are translated by 1 from their parents with probability $0 < p < 1$ and by -1 with probability $1 - p$. He proved that when $p > \frac{1}{2}$ the distribution of the centred maximum converges. But when $p = \frac{1}{2}$, Lifshits obtain a result analogous to (10), except

that the centring m_t is not explicit (it involves the median).

In the same way, Drmota [15], [18] and Chauvin and Drmota [9], proved Theorem 1.4 with analytical and combinatorial methods, again with an implicit m_t . We will see below that an explicit formula for m_t provides results like Corollary 1.2.

The average and the variance of the height and of the saturation level have been studied by many authors. Robson (1979) [30] found the asymptotic $\mathbb{E}(H_n) \asymp c^+ \log(n)$ and in 1995, Devroye and Reed [13] showed that $\text{Var}(H_n) = O_n((\log \log n)^2)$. These asymptotics have been improved [11], [12], [14], [27] until Reed [28] and Drmota [15] independently proved that $\mathbb{E}(H_n) = a_{\log(n)} + O_n(1)$ and that $\text{Var}(H_n) = O_n(1)$. Moreover for $n \in \mathbb{N}$, let $y_n(x)$ be recursively defined by:

$$\begin{cases} y_0(x) \equiv 1 \\ y_{n+1}(x) = 1 + \int_0^x y_n(t)^2 dt \end{cases} \quad (28)$$

Drmota showed that $y_n(1) = e^{\frac{n}{c^+} + \frac{3 \log n}{2(c^+ - 1)} + u_n + o_n(1)}$ where (u_n) is a bounded sequence such that $u_{n+1} - u_n \rightarrow 0$ and that if the better asymptotic

$$y_n(1) = e^{\frac{n}{c^+} + \frac{3 \log n}{2(c^+ - 1)} + D + o_n(1)}, \quad (29)$$

where $D \in \mathbb{R}$, is true then there are two 1-periodic continuous functions P_1 and P_2 such that:

$$\mathbb{E}(H_n) = a_{\log(n)} + P_1(a_{\log(n)}) + o_n(1), \quad (30)$$

$$\text{Var}(H_n) = P_2(a_{\log(n)}) + o_n(1). \quad (31)$$

Combining Drmota's result on the convergence of the distribution and Theorem 1.4, we get the following corollary.

Corollary 1.5. *The asymptotic (29) holds.*

In particular, this implies (30) and (31) which are the analogue results to Corollary 1.2. An interesting and open question is the asymptotic behaviour of the number of particles at the highest position. We will denote C_n this quantity for a binary search tree with n nodes and F_t for the Yule-time process at time t . By differentiating $\mathbb{E}(F_t)$ and applying to it Corollary 1.3 we obtain that: $\mathbb{E}(F_t) = P(a_t) + o_t(1)$ where P is a 1-periodic function. The fact that $y_n(1) = e^{\frac{n}{c^+} + \frac{3 \log n}{2(c^+ - 1)} + D + o_n(1)}$ and the work of Drmota [16] tells us that we have an analogue for $\mathbb{E}(C_n)$. Moreover, it is shown in [17] that the oscillations of $\mathbb{E}(C_n)$ around c^+ are at most of order 10^{-4} , which may suggest that $\mathbb{E}(C_n)$ converges. However it is quite possible to have small oscillations... This question is particularly interesting because it is related to a more general question, namely the convergence of the extremal process to a point process. Independently, Aïdekon, Berestycki, Brunet and Shi [3] and Arguin, Bovier and Kistler [5] have proved the convergence of the extremal process of branching Brownian motion and Madaule [26] of the branching random walk with non-lattice support. The lattice case has not been dealt with and the behaviour of $\mathbb{E}(F_t)$ could shed a first light on this case.

Finally, the question of the almost sure behaviour of the height and of the

saturation level for the binary search tree has been dealt with by Roberts in [29], where he showed that:

$$\frac{1}{2\theta^+} = \liminf_{n \rightarrow +\infty} \frac{c^+ \log(n) - H_n}{\log \log(n)} < \limsup_{n \rightarrow +\infty} \frac{c^+ \log(n) - H_n}{\log \log(n)} = \frac{3}{2\theta^+}, \quad (32)$$

and the analog result for the saturation level. We can derive Robert's result for the maximum and the minimum of the Yule-time process with ease.

2 Proof of Theorem 1.1

We recall that we obtain the branching random tree (T_n) by switching time and space in the process (X_t) (see (12)). The advantage of such a change of point of view, is that (T_n) is non-lattice and therefore Aïdekon's result [2] apply. As (T_n) and (X_t) are fundamentally the same object, this casts light on (X_t) as well and in particular provides the relation (13).

We further recall that the critical travelling-wave at speed c^+ has the following probabilistic representation:

$$\phi_K(x) = \mathbb{E} \left(e^{-K e^{-\theta^+ x} \partial W_\infty(\theta)} \right), \quad (33)$$

where $K > 0$ fixes the choice of the travelling wave, θ^+ is the unique solution in $\mathbb{R}^+$ of $c_\theta = c^+$ (see (8)) and $\partial W_\infty(\theta)$ is the limit of the derivative martingale defined by:

$$\partial W_t(\theta) = \text{sgn}(\theta) \sum_{u \in \mathcal{N}_t} (2e^\theta t \beta - X_u(t)) e^{\theta(X_u(t) - c_\theta t)}, \quad (34)$$

(it is a martingale with respect to $(\mathcal{F}_t)$, the filtration naturally associated to (X_t)). Moreover, we know [10] that the limit $\partial W_\infty(\theta)$ is almost surely positive for $\theta \in \{\theta^-, \theta^+\}$.

Proof. We want to prove that there exists $K > 0$ such that:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| = 0, \quad (35)$$

where $a_t = c^+ t - \frac{3 \log(t)}{2\theta^+}$. For $n \in \mathbb{N}$ and $x \in \mathbb{R}$, we define:

$$t_n = \frac{n}{c^+} + \frac{3 \log(n)}{2c^+ \theta^+} \quad \text{and} \quad g_n(x) = \mathbb{P} \left(T_{\min}(n) \geq t_n - \frac{x}{c^+} \right). \quad (36)$$

It can easily be seen that the branching random walk $(T_n)_{n \in \mathbb{N}}$ defined in (12) satisfies the hypothesis of Aïdekon's Theorem which yields the existence of a $K > 0$ such that:

$$g_n(x) \rightarrow \phi_K(x) \text{ uniformly in } x \quad (37)$$

(the uniformity follows from Dini's Theorem).

Thus, our strategy is to use (13) to prove (35) *via* Aïdekon's convergence result. We start by proving that there exists $K > 0$ such that for all $x \in \mathbb{R}$:

$$\lim_{t \rightarrow +\infty} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| = 0. \quad (38)$$

We first rewrite $\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor)$ by using (13):

$$\begin{aligned}
\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) &= \mathbb{P}(T_{\min}(\lfloor a_t + x \rfloor + 1) \geq t) \\
&= \mathbb{P}(T_{\min}(\lfloor a_t + x \rfloor + 1) \geq t_{\lfloor a_t + x \rfloor + 1} + t - t_{\lfloor a_t + x \rfloor + 1}) \\
&= \mathbb{P}\left(T_{\min}(\lfloor a_t + x \rfloor + 1) \geq t_{\lfloor a_t + x \rfloor + 1} - \frac{c^+(t_{\lfloor a_t + x \rfloor + 1} - t)}{c^+}\right) \\
&= g_{\lfloor a_t + x \rfloor + 1}(c^+(t_{\lfloor a_t + x \rfloor + 1} - t)) \tag{39}
\end{aligned}$$

A straightforward computation give:

$$c^+(t_{\lfloor a_t + x \rfloor + 1} - t) = \lfloor a_t + x \rfloor - a_t + 1 + \frac{3 \log(c^+)}{2\theta^+} + o_t(1),$$

where $o_t(1) = \frac{3}{2c^+\theta^+} \log\left(\frac{\lfloor a_t + x \rfloor + 1}{a_t + x + 1}\right)$ is a function that tends to 0 when t goes to infinity. By triangular inequality:

$$\begin{aligned}
&\left| g_{\lfloor a_t + x \rfloor + 1}(c^+(t_{\lfloor a_t + x \rfloor + 1} - t)) - \phi_K\left(\lfloor a_t + x \rfloor - a_t + 1 + \frac{3 \log(c^+)}{2\theta^+}\right) \right| \\
&\leq \left| g_{\lfloor a_t + x \rfloor + 1}(c^+(t_{\lfloor a_t + x \rfloor + 1} - t)) - \phi_K(c^+(t_{\lfloor a_t + x \rfloor + 1} - t)) \right| \\
&\quad + \left| \phi_K(c^+(t_{\lfloor a_t + x \rfloor + 1} - t)) - \phi_K\left(\lfloor a_t + x \rfloor - a_t + 1 + \frac{3 \log(c^+)}{2\theta^+}\right) \right|,
\end{aligned}$$

where K is fixed by (37). The first term in the right hand side goes to 0 by uniform convergence of (g_n) to ϕ_K (37) and the second goes to 0 by uniform continuity of ϕ_K (it is a continuous function with limits in $\pm\infty$).

We have thus proved that for all $x \in \mathbb{R}$:

$$\lim_{t \rightarrow +\infty} \left| \mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K\left(\lfloor a_t + x \rfloor - a_t + 1 + \frac{3 \log(c^+)}{2\theta^+}\right) \right| = 0,$$

which is (38) (the travelling-wave being defined up to an additive translation in the argument, we can incorporate $1 + \frac{3 \log(c^+)}{2\theta^+}$ into K).

Since the functions $x \mapsto \mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor)$ and $x \mapsto \phi_K(\lfloor a_t + x \rfloor - a_t)$ are both increasing and tends to 1 when x goes to $+\infty$ and to 0 when x goes to $-\infty$, for $\epsilon > 0$, there is $x_0 > 0$ such that there exists t_0 such that for all $t > t_0$:

$$\sup_{x \in]-\infty, -x_0] \cup [x_0, +\infty[} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| < \epsilon. \tag{40}$$

Moreover, since $[-x_0, x_0] \cap \mathbb{Z}$ is finite, we have that:

$$\lim_{t \rightarrow +\infty} \sup_{x \in [-x_0, x_0] \cap \mathbb{Z}} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| = 0. \tag{41}$$

For fixed t , $x \mapsto |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)|$ is constant on each interval of the form: $[\lfloor a_t \rfloor - a_t + k, \lfloor a_t \rfloor - a_t + k + 1[$, and then for fixed t

$$\begin{aligned}
&\sup_{x \in [-x_0, x_0] \cap \mathbb{Z}} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| \\
&= \sup_{x \in [-x_0, x_0]} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)|. \tag{42}
\end{aligned}$$

Combining (40), (41) and (42) we obtain:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(X_{\max}(t) \leq \lfloor a_t + x \rfloor) - \phi_K(\lfloor a_t + x \rfloor - a_t)| = 0. \quad (43)$$

By the same method, we recover the analogue result for the minimum. $\square$

We can now prove Corollaries 1.2 and 1.3.

3 Proof of Corollaries 1.2 and 1.3

Proof. To simplify, we denote ϕ instead of $\phi_{\max}$ the critical travelling-wave in Theorem 1.1. First, we need to establish estimates on the tails of the distribution of the maximum to determine a class of functions f sufficiently large for which H_f (14) and P_f (15) are well defined. Again, we will work with the Yule generation process. More precisely, we rely on a result due to Addario-Berry and Reed [1] which shows that the maximum of a branching random walk with i.i.d. displacements has exponential tails. Since our process does not have independent displacements, we can consider (T'_n) defined by:

$$T'_n = \frac{1}{2} \sum_{u \in \mathcal{M}_{n+1}} \delta_{T_u(n+1) - \tau_2}, \quad \forall n \geq 0, \quad (44)$$

where τ_2 is defined in (1). Note that (T'_n) is just a translation of (T_n) where we remove one element of each couple of particles. Since the relation (44) clearly implies:

$$T_{\min}(n) = T'_{\min}(n-1) + \tau_2 \quad (45)$$

and by using Addario-Berry and Reed's result [1, Theorem 3] we obtain by straightforward computations that there is $C > 0$ and $\alpha > 0$ such that for all $n \geq 1$ and $x \in \mathbb{R}^+$:

$$\mathbb{P}(|T_{\min}(n) - t_n| \geq x) \leq C e^{-\alpha x}, \quad (46)$$

where t_n is defined by (36). Again, using (13) and taking $n = \lfloor a_t \rfloor + k$ in (46), we obtain that there exist $C' > 0$ and $\alpha' > 0$ such that for all $t \geq 1$ and all $k \in \mathbb{N}$,

$$\mathbb{P}(|X_{\max}(t) - \lfloor a_t \rfloor| \geq k) \leq C' e^{-\alpha' k}. \quad (47)$$

Moreover, we precisely know the asymptotics of ϕ by [8] and (27),

$$1 - \phi(x) \underset{x \rightarrow +\infty}{\sim} x e^{-\theta x} \quad \text{and} \quad \phi(x) \underset{x \rightarrow -\infty}{\sim} e^{\frac{\beta x}{c^+}}. \quad (48)$$

Let $\delta > 0$, such that $\delta < \min\left(\theta, \frac{\beta}{c^+}, \alpha'\right)$. Let f be a piecewise continuous function, such that $f(x) = o_{\pm\infty}(e^{\delta|x|})$ and for $k \in \mathbb{Z}$, let h_k and p_k be defined by:

$$\begin{aligned} h_k(t) &= f(k - a_t) \mathbb{P}(X_{\max}(t) = k) \\ &= f(k - a_t) [\mathbb{P}(X_{\max}(t) \leq k) - \mathbb{P}(X_{\max}(t) \leq k-1)], \quad \forall t > 0 \end{aligned} \quad (49)$$

and

$$p_k(s) = f(k - s) [\phi(k - s) - \phi(k - 1 - s)], \quad \forall s \in \mathbb{R}. \quad (50)$$

The asymptotics in (47), (48) and the assumptions on f imply that there exists $(\delta', A) \in (\mathbb{R}^*)^2$ such that for all $(k, t) \in \mathbb{Z} \times [1, +\infty)$:

$$h_{k+\lfloor a_t \rfloor}(t) \leq Ae^{-\delta'|k|} \quad \text{and} \quad p_{k+\lfloor a_t \rfloor}(a_t) \leq Ae^{-\delta'|k|}. \quad (51)$$

We see that the sums P_f and H_f are invariant by translation by an integer. Thus:

$$H_f(t) = \sum_{k \in \mathbb{Z}} h_{k+\lfloor a_t \rfloor}(t) \quad \text{and} \quad P_f(a_t) = \sum_{k \in \mathbb{Z}} p_{k+\lfloor a_t \rfloor}(a_t) \quad (52)$$

The upper bound (51) tell us that $H_f(t)$ and $P_f(a_t)$ are well-defined and that for $\epsilon > 0$, there exist $k_0 \in \mathbb{Z}^-$ and $k_1 \in \mathbb{N}$ such that for all $t \geq 1$:

$$\sum_{k=-\infty}^{k_0} h_{k+\lfloor a_t \rfloor}(t) + \sum_{k=k_1}^{+\infty} h_{k+\lfloor a_t \rfloor}(t) < \epsilon, \quad (53)$$

and

$$\sum_{k=-\infty}^{k=k_0} p_{k+\lfloor a_t \rfloor}(a_t) + \sum_{k=k_1}^{+\infty} p_{k+\lfloor a_t \rfloor}(a_t) < \epsilon. \quad (54)$$

Furthermore, Theorem 1.1 ensures that :

$$\lim_{t \rightarrow +\infty} \sum_{k=k_0}^{k=k_1} |h_{k+\lfloor a_t \rfloor}(t) - p_{k+\lfloor a_t \rfloor}(a_t)| = 0. \quad (55)$$

Corollary 1.2 is thus proved. We have yet to deal with Corollary 1.3. For $m \in \mathbb{N}$, let $f \in \mathcal{C}^m(\mathbb{R})$ with:

$$f^{(i)}(x) = o_{\pm\infty}(e^{\delta|x|}), \quad \forall 0 \leq i \leq m. \quad (56)$$

This obviously implies that: $t \mapsto p_k(a_t)$ and h_k belong to $\mathcal{C}^m(\mathbb{R}_+^*)$. Let $h(x, t) = \mathbb{P}(X_{\max}(t) \leq x)$, for $(x, t) \in \mathbb{N} \times \mathbb{R}_+^*$ and let $i \in [1, m]$. If we differentiate $i - 1$ times equation (5) we get:

$$\partial_t^{(i)} h(x, t) = \beta \left(\sum_{j=0}^{i-1} \binom{i-1}{j} \partial_t^{(j)} h(x-1, t) \partial_t^{(i-j-1)} h(x-1, t) - \partial_t^{(i-1)} h(x, t) \right). \quad (57)$$

By an immediate induction, there is $C_i > 0$, such that for x large enough and $t > 0$:

$$\partial_t^{(i)} h(x, t) \leq C_i e^{-\alpha'(k-\lfloor a_t \rfloor)}. \quad (58)$$

Let K be a compact set of $\mathbb{R}_+^*$. $\lfloor a_t \rfloor$ is bounded on K and therefore the hypothesis (56) on f and equation (58) gives us the existence of C'_i such that we have:

$$h_k^{(i)}(t) \leq C'_i e^{-(\alpha'-\delta)k}, \quad \forall t \in K. \quad (59)$$

The series $(\sum h_k^{(i)})$ then converges uniformly on every compact of $\mathbb{R}_+^*$ for each $i \in [0, m]$, which implies that $H_f \in \mathcal{C}^m(\mathbb{R}_+^*)$. With the same approach, we can show that $P_f \in \mathcal{C}^m(\mathbb{R})$. To prove the final part of the corollary, we can again

cut the sum in three terms and proceed as for the previous corollary. We will simply show by induction that:

$$\lim_{t \rightarrow +\infty} \sup_{x \in \mathbb{Z}} |\partial_t^{(i)} h(x + \lfloor a_t \rfloor, t) - (-c)^i \phi^{(i)}(x - \{a_t\})| = 0 \quad (60)$$

The base case $i = 0$ is Theorem 1.1. If we assume that the result holds for the rank $i - 1$, we can rewrite (57):

$$\begin{aligned} \frac{\partial_t^{(i)} h(x, t)}{(-c)^{i-1} \beta} &= \sum_{j=0}^{i-1} \binom{i-1}{j} \phi^{(j)}(x-1-\{a_t\}) \phi^{(j-i+1)}(x-1-\{a_t\}) \\ &\quad - \phi^{(i-1)}(x - \{a_t\}) + o_t(1), \end{aligned} \quad (61)$$

where $o_t(1)$ is uniform in x . Differentiating $i - 1$ times equation (6) and combining the result with (61), we obtain (60).

The end of the proof is identical as for Corollary 1.2. $\square$

For the minimum the proof is the same except that one of the asymptotics of $\phi_{\min}$ (48) is different from those of $\phi_{\max}$. Indeed, $\phi_{\min}$ is decreasing and thus (6) implies that for all $x \in \mathbb{R}$, $\phi_{\min}(x) \leq \phi_{\min}^2(x-1)$. Consequently, there is $0 < A < 1$ such that:

$$\phi_{\min}(x) \leq A^{2^x}, \quad \forall x > 0.$$

4 Application to the binary search tree

We now know the asymptotic behaviour of the maximum of the Yule-time process. Thanks to that, and using the connection between the Yule process and the binary search tree, we will be able to prove Theorem 1.4.

Recall that the derivative martingale of the Yule tree (34) is the derivative with respect to θ (up to a change of sign for $\theta < 0$) of the additive martingale defined as:

$$W_t(\theta) = \sum_{u \in \mathcal{N}_t} e^{\theta(X_u(t) - c_\theta t)}. \quad (62)$$

We know [10] that this martingale converges to an almost surely positive random variable when $\theta \in]\theta^-, \theta^+[$, and in particular that the limit of the martingale is an exponential random variable with parameter 1 when $\theta = 0$.

Likewise, for the binary search tree, let $\mathcal{L}_n$ be the set of leaves of $\mathcal{T}_n$ and define $\mathcal{F}_n^{BST} = \sigma(\{u \in \mathcal{T}_i\}, i \leq n, u \in \mathcal{T})$ where $\mathcal{T}$ is the complete binary tree. For $z \in \mathbb{R} \setminus -\frac{1}{2}\mathbb{N}$, the process:

$$M_n(z) = \frac{1}{C_n(z)} \sum_{u \in \mathcal{L}_n} z^{|u|}, \quad (63)$$

where:

$$C_0(z) = 1 \quad \text{and} \quad C_n(z) = \prod_{k=0}^{n-1} \frac{k+2z}{k+1} \quad (64)$$

is a $(\mathcal{F}_n^{BST})$ -martingale introduced by Jabbour [20], which we will call the additive martingale of the binary search tree. As for the Yule-time process, we can

define the derivative martingale of the binary search tree:

$$\partial M_n(z) = \frac{dM_n(z)}{dz}. \quad (65)$$

We now recall that we can embed the binary search tree into the Yule tree (2). For $n \geq 2$, let τ_n be the time $\inf\{t > 0, N_t = n\}$ and $\tau_1 = 0$. As mentioned above if we stop the Yule tree process at time τ_n , the tree we obtain is a random binary search tree. We call Consequently if H_n is the height of a random binary search tree embedded into the Yule process, we have

$$H_n = X_{\max}(\tau_n). \quad (66)$$

The following theorem proved by Chauvin and Rouault give us more information on the derivative martingale (65) and on its connection with the Yule process.

Theorem (Chauvin, Rouault [8]). *The martingale $(\partial M_n(z))$ converges as $n \rightarrow +\infty$ to a positive random variable ∂M_∞^+ for $z = e^{\theta^+}$ (resp. to ∂M_∞^- for $z = e^{\theta^-}$). Moreover, by embedding the binary search tree into the Yule process, we have:*

$$\partial W_\infty(\theta^+) = \frac{e^{-\theta^+} W_\infty(0)^{c^++1}}{\Gamma(c^+)} \partial M_\infty^+ \quad (67)$$

and:

$$\partial W_\infty(\theta^-) = \frac{e^{-\theta^-} W_\infty(0)^{c^-+1}}{\Gamma(c^-)} \partial M_\infty^-, \quad (68)$$

where the limit of the additive martingale $W_\infty(0)$ has a random exponential 1 law and is independent of ∂M_∞^+ and ∂M_∞^- .

Drawing our inspiration from Lalley and Selke [22] and with the help of the former theorem, we will prove Theorem 1.4.

Proof. For $x \in \mathbb{R}$, recall that $\{x\} := x - \lfloor x \rfloor$

Step 1: First, we will show that:

$$\exists K > 0 : \forall \epsilon > 0 \exists s_0 > 0 : \forall s > s_0 \exists t_s > 0 : \forall t > t_s :$$

$$\left| \mathbb{P}(X_{\max}(t + s - \log(W_s)) \leq \lfloor a_{t+s} + x \rfloor) - \mathbb{E} \left(\exp \left(-K e^{-\theta(\lfloor a_{t+s} + x \rfloor - a_{t+s})} \partial M_\infty(\theta) \right) \right) \right| < \frac{\epsilon}{3}, \quad (69)$$

where $W_s = W_s(0)$. First, by Markov property:

$$\begin{aligned} & \mathbb{P}(X_{\max}(t + s - \log(W_s)) \leq a_{t+s} + x | \mathcal{F}_s) \\ &= \prod_{u \in \mathcal{N}_s} \mathbb{P}(X_{\max}^u(t - \log(W_s)) \leq a_{t+s} + x - X_u(s)) \\ &= \prod_{u \in \mathcal{N}_s} \mathbb{P}(X_{\max}^u(t - \log(W_s)) \leq a_{t-\log W_s} + a_{t+s} + x - X_u(s) - a_{t-\log W_s}), \end{aligned} \quad (70)$$

where the $X_{\max}^u$ are independent of $\mathcal{F}_s$ and identically distributed. For a fixed s we have by Theorem 1.1 that almost surely:

$$\lim_{t \rightarrow +\infty} \left| \mathbb{P} \left(X_{\max}(t + s - \log(W_s)) \leq a_{t+s} + x \mid \mathcal{F}_s \right) - \prod_{u \in \mathcal{N}_s} \phi_K(\lfloor a_{t+s} + x \rfloor - a_{t-\log W_s} - X_u(s)) \right| = 0, \quad (71)$$

where $\phi_K(x) = \mathbb{E} \left(\exp \left(-K e^{-\theta^+ x} \partial W_{\infty}(\theta^+) \right) \right)$. Observe that the argument of ϕ_K can be rewritten as:

$$\begin{aligned} \lfloor a_{t+s} + x \rfloor - a_{t-\log W_s} - X_u(s) &= a_{t+s} + x - \{a_{t+s} + x\} - a_{t-\log(W_s)} - X_u(s) \\ &= c^+ s + c^+ \log W_s + R_{x,s,t} + S_{s,t} - X_u(s), \end{aligned}$$

where

$$R_{x,s,t} = x - \{a_{t+s} + x\} \quad \text{and} \quad S_{s,t} = \frac{3 \log(\frac{t - \log W_s}{t+s})}{2\theta^+}.$$

Moreover, $S_{s,t}$ goes to 0 when t goes to infinity. ϕ_K is continuous and bounded by 1, therefore we have by dominated convergence:

$$\begin{aligned} \lim_{t \rightarrow +\infty} \left| \mathbb{P} \left(X_{\max}(t + s - \log(W_s)) \leq a_{t+s} + x \right) - \mathbb{E} \left(\prod_{u \in \mathcal{N}_s} \phi_K(c^+ s + c^+ \log W_s + R_{x,s,t} - X_u(s)) \right) \right| &= 0 \end{aligned} \quad (72)$$

We will now rewrite the product of the terms from (72):

$$\prod_{u \in \mathcal{N}_s} \phi_K(c^+ s + c^+ \log W_s + R_{x,s,t} - X_u(s)) \quad (73)$$

$$= \prod_{u \in \mathcal{N}_s} \mathbb{E} \left(\exp \left(-K e^{-\theta(c^+ s + c^+ \log W_s + R_{x,s,t} - X_u(s))} \partial W_{\infty,u}(\theta^+) \right) \mid \mathcal{F}_s \right) \quad (74)$$

$$= \mathbb{E} \left(\exp \left(-K e^{-\theta(c^+ \log W_s + R_{x,s,t})} \sum_{u \in \mathcal{N}_s} e^{\theta(X_u(s) - c^+ s)} \partial W_{\infty,u}(\theta^+) \right) \mid \mathcal{F}_s \right), \quad (75)$$

where $\partial W_{\infty,u}(\theta^+)$ are independent copies of $\partial W_{\infty}(\theta^+)$ and independent of $\mathcal{F}_s$. Moreover we know that:

$$\sum_{u \in \mathcal{N}_s} e^{\theta(X_u(s) - c^+ s)} \partial W_{\infty,u}(\theta^+) = \partial W_{\infty}(\theta^+) = \frac{W_{\infty}(0)^{c^++1}}{\Gamma(c^+)} \partial M_{\infty}(\theta^+). \quad (76)$$

The definition of c^+ gives us $c^+ \theta^+ = c^+ + 1$ and thus:

$$\begin{aligned} &\prod_{u \in \mathcal{N}_s} \phi_K(c^+ s - X_u(s) + c^+ \log W_s + R_{x,s,t}) \\ &= \mathbb{E} \left(\exp \left(-K e^{-\theta^+(c^+ \log W_s + R_{x,s,t})} \frac{W_{\infty}(0)^{c^++1}}{\Gamma(c^+)} \partial M_{\infty}(\theta^+) \right) \mid \mathcal{F}_s \right) \\ &= \mathbb{E} \left(\exp \left(-K' e^{-\theta^+ R_{x,s,t}} \left(\frac{W_{\infty}(0)}{W_s(0)} \right)^{c^++1} \partial M_{\infty}(\theta^+) \right) \mid \mathcal{F}_s \right), \end{aligned} \quad (77)$$

with $K' = \frac{K}{\Gamma(c^+)}$. Let $\Psi_{K'}(R, s) = \mathbb{E} \left(\exp \left(-K' e^{-\theta^+ R} \left(\frac{W_\infty(0)}{W_s(0)} \right)^{c^++1} \partial M_\infty(\theta^+) \right) \right)$.

By dominated convergence:

$$\lim_{s \rightarrow +\infty} \Psi_{K'}(R, s) = \psi(R).$$

Moreover for fixed s , $\Psi(\cdot, s)$ is increasing and

$$\lim_{x \rightarrow +\infty} \Psi(x, s) = 1 \quad \text{and} \quad \lim_{x \rightarrow -\infty} \Psi(x, s) = 0.$$

Thus by Dini's Theorem:

$$\lim_{s \rightarrow +\infty} \sup_{R \in \mathbb{R}} |\Psi_{K'}(R, s) - \psi(R)| = 0.$$

We then have:

$$\lim_{s \rightarrow +\infty} \sup_{t > 0} \left| \mathbb{E} \left(\prod_{u \in \mathcal{N}_s} \phi_K(c^+ s - X_u(s) + c^+ \log W_s + R_{x,s,t}) \right) - \psi_{K'}(R_{x,s,t}) \right| = 0. \quad (78)$$

So from equations (72) and (78) we deduce (69).

Step 2: Noticing that $\log(n) - \log(W_{\tau_n}(0)) = \tau_n$, we have by (66):

$$\begin{aligned} \mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) &= \mathbb{P}(X_{\max}(\tau_n) \leq \lfloor a_{\log(n)} + x \rfloor) \\ &= \mathbb{P}(X_{\max}(\log(n) - \log(W_{\tau_n}(0))) \leq \lfloor a_{\log(n)} + x \rfloor). \end{aligned} \quad (79)$$

For every $K > 0$, ψ_K is uniformly continuous and hence for $\epsilon > 0$ there is $\eta > 0$ such that for all $(x, y) \in \mathbb{R}$ such that $|x - y| < \eta$,

$$|\psi_K(x) - \psi_K(y)| < \frac{\epsilon}{3}. \quad (80)$$

For $\eta' > 0$, we may choose $s_1 > s_0$ (where s_0 is defined in (69)) such that for every $s > s_1$:

$$\mathbb{P}(|\log(W_s(0)) - \log(W_\infty(0))| > \frac{\eta'}{2}) < \frac{\epsilon}{6}. \quad (81)$$

by the almost sure convergence of $W_s(0)$ to a positive random variable. In the same way, there exists $n_1 \in \mathbb{N}$ such that: $\log(n_1) > t_{s_1} + s_1$ and such that for all $n \geq n_1$:

$$\mathbb{P}(|\log(W_{\tau_n}(0)) - \log(W_\infty(0))| > \frac{\eta'}{2}) < \frac{\epsilon}{6}. \quad (82)$$

Introducing the approximations (81) and (82) in the equation (79) and using the monotonicity of $\psi_{K'}$ (where K' is defined in (77)) we obtain for $n \geq n_1$:

$$\begin{aligned} \mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) &\leq \mathbb{P}(X_{\max}(\log(n) - \log(W_{s_1}(0)) - \eta') \leq \lfloor a_{\log(n)} + x \rfloor) + \frac{\epsilon}{3} \\ &\leq \psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n) - \eta'}) + \frac{2\epsilon}{3}, \end{aligned} \quad (83)$$

where equation (83) is a consequence of (69). So, taking $\eta' > 0$ such that for all $n \geq n_1$, $|a_{\log(n) - \eta'} - a_{\log(n)}| < \eta$, we obtain:

$$\begin{aligned} \mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) &\leq \psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n)} + \eta) + \frac{2\epsilon}{3} \\ &\leq \psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n)}) + \epsilon, \end{aligned} \quad (84)$$

by (80). Similarly, we obtain:

$$\psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n)}) - \epsilon \leq \mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor), \quad (85)$$

and then for $x \in \mathbb{R}$,

$$\lim_{n \rightarrow +\infty} |\mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) - \psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n)})| = 0.$$

By the same arguments as in the proof of Theorem 1.1, we have in fact that:

$$\lim_{n \rightarrow +\infty} \sup_{x \in \mathbb{R}} |\mathbb{P}(H_n \leq \lfloor a_{\log(n)} + x \rfloor) - \psi_{K'}(\lfloor a_{\log(n)} + x \rfloor - a_{\log(n)})| = 0. \quad (86)$$

□

Acknowledgements

I am very grateful to my thesis advisor, Julien Berestycki, for his help throughout the writing of this article. I would also thank Joon Kwon for very useful comments.

References

- [1] Louigi Addario-Berry and Bruce Reed. Minima in branching random walks. *The Annals of Probability*, pages 1044–1079, 2009.
- [2] Elie Aïdékon. Convergence in law of the minimum of a branching random walk. *The Annals of Probability*, 41(3A):1362–1426, 2013.
- [3] Elie Aïdékon, Julien Berestycki, Éric Brunet, and Zhan Shi. Branching brownian motion seen from its tip. *Probability Theory and Related Fields*, 157(1-2):405–451, 2013.
- [4] Gerold Alsmeyer, JD Biggins, Matthias Meiners, et al. The functional equation of the smoothing transform. *The Annals of Probability*, 40(5):2069–2105, 2012.
- [5] Louis-Pierre Arguin, Anton Bovier, and Nicola Kistler. The extremal process of branching brownian motion. *Probability Theory and Related Fields*, 157(3-4):535–574, 2013.
- [6] Maury Bramson. *Convergence of solutions of the Kolmogorov equation to travelling waves*, volume 44. American Mathematical Society, 1983.
- [7] Maury Bramson, Jian Ding, and Ofer Zeitouni. Convergence in law of the maximum of nonlattice branching random walk. *arXiv preprint arXiv:1404.3423*, 2014.
- [8] B. Chauvin and A. Rouault. Connecting yule process, bisection and binary search tree via martingales. *JIRSS*, 3(2):88–116, 2004.
- [9] Brigitte Chauvin and Michael Drmota. The random multisection problem, travelling waves and the distribution of the height of m-ary search trees. *Algorithmica*, 46(3-4):299–327, 2006.

- [10] Brigitte Chauvin, Thierry Klein, Jean-Francois Marckert, and Alain Rouault. Martingales and profile of binary search trees. *Electron. J. Probab*, 10(12):420–435, 2005.
- [11] Luc Devroye. A note on the height of binary search trees. *Journal of the ACM (JACM)*, 33(3):489–498, 1986.
- [12] Luc Devroye. Branching processes in the analysis of the heights of trees. *Acta Informatica*, 24(3):277–298, 1987.
- [13] Luc Devroye and Bruce Reed. On the variance of the height of random binary search trees. *SIAM Journal on Computing*, 24(6):1157–1162, 1995.
- [14] Michael Drmota. An analytic approach to the height of binary search trees. *Algorithmica*, 29(1-2):89–119, 2001.
- [15] Michael Drmota. An analytic approach to the height of binary search trees ii. *Journal of the ACM (JACM)*, 50(3):333–374, 2003.
- [16] Michael Drmota. On robson’s convergence and boundedness conjectures concerning the height of binary search trees. *Theoretical computer science*, 329(1):47–70, 2004.
- [17] Michael Drmota. Profile and height of random binary search trees. *J. Iranian Statistical Society*, 3(2):117–138, 2004.
- [18] Michael Drmota. *Random trees: an interplay between combinatorics and probability*. Springer, 2009.
- [19] Simon C Harris. Travelling-waves for the fkpp equation via probabilistic arguments. *Proceedings of the Royal Society of Edinburgh: Section A Mathematics*, 129(03):503–517, 1999.
- [20] Jean Jabbour-Hattab. Martingales and large deviations for binary search trees. *Random Structures & Algorithms*, 19(2):112–127, 2001.
- [21] AN Kolmogorov, IG Petrovskii, and NS Piskunov. A study of the equation of diffusion with increase in the quantity of matter, and its application to a biological problem. *Bjul. Moskovskogo Gos. Univ*, 1(7):1–26, 1937.
- [22] SP Lalley, T Sellke, et al. A conditional limit theorem for the frontier of a branching brownian motion. *The Annals of Probability*, 15(3):1052–1061, 1987.
- [23] Mikhail Anatolievich Lifshits. Cyclic behavior of maxima in a hierarchical summation scheme. *arXiv preprint arXiv:1212.0189*, 2012.
- [24] Quansheng Liu. Fixed points of a generalized smoothing transformation and applications to the branching random walk. *Advances in Applied Probability*, pages 85–112, 1998.
- [25] Quansheng Liu. On generalized multiplicative cascades. *Stochastic processes and their applications*, 86(2):263–286, 2000.
- [26] Thomas Madaule. Convergence in law for the branching random walk seen from its tip. *arXiv preprint arXiv:1107.2543*, 2011.

- [27] Boris Pittel. On growing random binary trees. *Journal of Mathematical Analysis and Applications*, 103(2):461–480, 1984.
- [28] Bruce Reed. The height of a random binary search tree. *Journal of the ACM (JACM)*, 50(3):306–332, 2003.
- [29] Matthew I Roberts. Almost sure asymptotics for the random binary search tree. *arXiv preprint arXiv:1002.3896*, 2010.
- [30] John Michael Robson. The height of binary search trees. *Australian Computer Journal*, 11(4):151–153, 1979.